


Today

- Sparsest cut
- Reduction to H-Sparsest cut
- $O(\log n)$ -approximation for H-sparsest cut via Bourgain + cut cone
- Expanders



Recall


Given $x, y \in \mathbb{R}^n$ define $d_1(x, y) := \sum_i |x_i - y_i|$

Bourgain's Theorem: given any n -point metric (V, δ) , \exists (poly-time computable) embedding f w/ distortion $O(\log n)$ of (V, δ) into (\tilde{V}, d_1) for $\tilde{V} \subseteq \mathbb{R}^{O(\log^2 n)}$

1-1 Correspondence Between Metrics and Vectors


$$(V, d) \quad \text{s.t. } V = \{v_1, v_2, \dots, v_n\} \quad \leftrightarrow \quad \begin{pmatrix} d(v_1, v_2) \\ d(v_1, v_3) \\ \vdots \\ d(v_1, v_n) \\ d(v_2, v_3) \\ \vdots \end{pmatrix} \in \mathbb{R}^{\binom{n}{2}}$$

(V, δ) is an ℓ_1 metric if it embeds isometrically into (\tilde{V}, d_1) for $\tilde{V} \subseteq \mathbb{R}^n$ for some n




(V, d_S) is a cut metric for $S \subseteq V$ s.t.

$$d_S(u, v) = \begin{cases} 0 & u, v \in S \text{ or } u, v \notin S \\ 1 & \text{otherwise} \end{cases}$$



Let $\mathcal{L}_1(V) \subseteq \mathbb{R}^{\binom{n}{2}}$ be all ℓ_1 metrics on V and let $\text{Cut}(V) \subseteq \mathbb{R}^{\binom{n}{2}}$ be all cut metrics of V

The convex cone of $D \subseteq \mathbb{R}^{\binom{n}{2}}$ is $\text{Cone}(D) := \left\{ \sum_{d \in D} \alpha_d \cdot d : \alpha_d \geq 0 \forall d \right\}$



Theorem: $\mathcal{L}_1(V) = \text{Cone}(\text{Cut}(V)) \quad \forall V.$

Also, given $d \in \mathcal{L}_1(V)$, \exists ℓ_1 embedding f can poly-time compute $d_i \in \text{Cut}(V), \alpha_i \in \mathbb{R} \forall i$ s.t. $d = \sum_i \alpha_i \cdot d_i$

Sparsest Cut

Given connected graph $G=(V,E)$

The volume of $S \subseteq V$ is $Vol(S) := \sum_{u \in S} deg(u)$


The conductance of $S \subseteq V$ is $\Phi(S) := \frac{|\delta(S)|}{\min(Vol(S), Vol(V \setminus S))}$

Sparsest Cut Problem: find non-empty $S \subseteq V$ minimizing $\Phi(S)$

Let $OPT := \min_S \Phi(S)$

Note: not always a min s-t cut

Note: $\Phi(S) \in (0, 1]$ always b/c $\min(Vol(S), Vol(V \setminus S)) \geq |\delta(S)|$



Fact: Sparsest Cut is NP-hard


H-Sparsest Cut

$w: E \rightarrow \mathbb{R}_{\geq 0}$

Also given complete edge-weighted graph $H=(V, E_H, w)$

The H-conductance of $S \subseteq V$ is $\Phi_H(S) := \frac{|\delta_G(S)|}{w(\delta_H(S))} \rightarrow \sum_{e \in \delta_H(S)} w(e)$

H-Sparsest Cut Problem: find non-empty $S \subseteq V$ minimizing $\Phi_H(S)$



Theorem: \exists Poly-time $O(\log n)$ -approximation for H-Sparsest Cut

Corollary: \exists Poly-time $O(\log n)$ -approximation for Sparsest Cut

Let $w_H(\{u,v\}) = \frac{deg_G(u) \cdot deg_G(v)}{m}$; will show $\Phi(S) \leq \Phi_H(S) \leq 2 \cdot \Phi(S) \quad \forall S \subseteq V$ → Corollary follows b/c objective changed by ≤ 2

$\forall u,v$ and let $V_S = Vol(S), \bar{V}_S = Vol(V \setminus S)$

so $w(\delta_H(S)) = \sum_{\substack{u \in S \\ v \in \bar{S}}} \frac{deg_G(u) \cdot deg_G(v)}{m} = \frac{1}{m} \sum_{u \in S} deg_G(u) \sum_{v \in \bar{S}} deg_G(v) = \frac{1}{m} \sum_{u \in S} deg_G(u) \cdot \bar{V}_S = \frac{1}{m} V_S \cdot \bar{V}_S$

$\frac{m}{2} \cdot \min(V_S, \bar{V}_S) \leq \min(V_S, \bar{V}_S) \cdot \max(V_S, \bar{V}_S) = V_S \cdot \bar{V}_S \leq m \cdot \min(V_S, \bar{V}_S)$

so $w(\delta_H(S)) \in [\frac{1}{2} \cdot \min(V_S, \bar{V}_S), \min(V_S, \bar{V}_S)]$

so $\forall S \subseteq V \quad \Phi(S) \leq \Phi_H(S) \leq 2 \cdot \Phi(S)$

H-Sparsest Cut via Cut Metrics

Variable $d(u,v)$ for each $u,v \in V$; let $d(e) := d(u,v)$ for $e = (u,v)$

$$\min_d \frac{\sum_{e \in E} d(e)}{\sum_{u,v} d(u,v) \cdot w(e_{u,v})} \quad \text{s.t.} \quad d \text{ is a cut metric on } V \quad (1)$$

Not an LP !!

↓ Leighton-Rao Relaxation

$$\min_d \frac{\sum_{e \in E} d(e)}{\sum_{u,v} d(u,v) \cdot w(e_{u,v})} \quad \text{s.t.} \quad \begin{cases} d(x,x) = 0 \quad \forall x \\ d(x,y) = d(y,x) \quad \forall x,y \\ d(x,y) \leq d(x,z) + d(z,y) \quad \forall x,y,z \end{cases} \quad (2)$$

"d is a metric"

↕

$$\min_d \frac{\sum_{e \in E} d(e)}{\sum_{u,v} d(u,v) \cdot w(e_{u,v})} \quad \text{s.t.} \quad \begin{cases} d(x,x) = 0 \quad \forall x \\ d(x,y) = d(y,x) \quad \forall x,y \\ d(x,y) \leq d(x,z) + d(z,y) \quad \forall x,y,z \\ \sum_{u,v} d(u,v) \cdot w(e_{u,v}) = 1 \end{cases} \quad (3)$$

An LP !!

Let OPT_{LR} be optimal value of

Claim: $OPT_{LR} \leq OPT$

Let OPT_i be optimal of (i) above and O_i be objective of (i)

$OPT_2 \leq OPT_1 = OPT$

↑ min over larger set
↑ usual 1-1 corr.

$OPT_{LR} = OPT_3 \leq OPT_2$

$OPT_3 \leq OPT_2$ [Suppose d optimal for (2) and let $D = \sum_{u,v} d(u,v) \cdot w(e_{u,v})$ $OPT_2 = O_2(d) = \frac{O_3(d)}{D}$
 But $\frac{1}{D} \cdot d$ is feasible for (3) so $OPT_3 \leq O_3(\frac{1}{D} \cdot d) = \frac{O_3(d)}{D} = OPT_2$

$OPT_2 \leq OPT_3$ [Suppose d_3 optimal for (3) so $OPT_3 = O_3(d_3)$ extra constraint of (2)
 But d_3 feasible for (2) so $OPT_2 \leq O_2(d_3) = O_3(d_3) = OPT_3$

Proof of Theorem

Idea: Solution to (3) $\xrightarrow{\text{Bourgain}} \ell_1$ metric $\xrightarrow{\text{Cut Cone}} \text{Cut on } G$
 lose $O(\log n)$

Alg

Let d be an optimal solution to (3)

Let (\tilde{v}, d) be what Bourgain embeds (V, d) into w/ $O(\log n)$ distortion

$d_1 \in \text{Cone}(\text{cut}(\tilde{v}))$ so $d_1 = \sum_{S \in \mathcal{L}} \alpha_S \cdot d_S$ s.t. $\alpha_S \geq 0$ and d_S a cut metric w/ cut S

Let S^* be $S \in \mathcal{L}$ minimizing $\Phi_H(S)$

Return S^*

Poly-time b/c Bourgain + Cut Cone poly-time doable

To see approximation guarantee:

For $S \in \mathcal{L}$ let $N_S := \sum_e d_S(e)$ and $D_S := \sum_{u,v} d_S(u,v) \cdot w(u,v)$

$$\text{So } S^* = \underset{S \in \mathcal{L}}{\text{argmin}} \frac{N_S}{D_S} = \underset{S \in \mathcal{L}}{\text{argmin}} \frac{\alpha_S \cdot N_S}{\alpha_S \cdot D_S}$$

$$\Phi_H(S^*) = \frac{N_{S^*}}{D_{S^*}} = \frac{\alpha_{S^*} \cdot N_{S^*}}{\alpha_{S^*} \cdot D_{S^*}} \leq \frac{\sum_S \alpha_S \cdot N_S}{\sum_S \alpha_S \cdot D_S} = \frac{\sum_e \sum_S \alpha_S d_S(e)}{\sum_{u,v} w(u,v) \sum_S \alpha_S d_S(u,v)} = \frac{\sum_e d_1(e)}{\sum_{u,v} w(u,v) \cdot d_1(u,v)}$$

$$\min_i \frac{a_i}{b_i} \leq \frac{\sum_i a_i}{\sum_i b_i}$$



$$\leq O(\log n) \cdot \frac{\sum_e d(e)}{\sum_{u,v} d(u,v) \cdot w(u,v)} = O(\log n) \cdot \text{OPT}_{LR} \leq O(\log n) \cdot \text{OPT}$$

↑
↑
Bourgain has $O(\log n)$ distortion
claim

Expanders

$G=(V,E)$ is a ϕ -edge-expander if $\phi(S) \geq \phi \quad \forall$ non-empty $S \subset V$

E.g.



① K_n is an $\Omega(1)$ -expander

② Above is a $O(\frac{1}{n^2})$ -expander


Complete graph on n vertices

Among most well-studied graph classes in TCS. Why?

Useful for impossibility results

$\deg(v) = 3 \forall v$

Fact: $\forall n_0, \exists$ an $n \geq n_0$ -node 3-regular $\Omega(1)$ -expander




\exists sparse graphs that are as connected as K_n

Shows:

- LDD separation quality $\Omega(\frac{\log n}{n})$
- Embeddings into l_1 or l_2 require $\Omega(\log n)$ distortion (tightness of Bourgain)
- "Probabilistic tree embeddings" require $\Omega(\log n)$ distortion
- Many others...

Useful for algorithms

A ϕ -expander decomposition of $G=(V,E)$ is $F \subseteq E$ s.t. that the connected components of $(V, E \setminus F)$ are ϕ -expanders



Fact: \forall m -edge graphs $\forall \phi \in (0,1), \exists$ a ϕ -expander decomposition F s.t. $|F| \leq O(\phi \cdot \log n \cdot m)$

Fact: Poly-time α -approximate sparsest cut \rightarrow Poly-time ϕ -expander decomposition. F s.t. $|F| \leq O(\alpha \cdot \phi \cdot \log n \cdot m)$